

- Lassen, J., and Løvendahl, P. 2016. Heritability estimates for enteric methane emissions from Holstein cattle measured using noninvasive methods. *J. Dairy Sci.* 99 (3), 1959-1967.
<https://doi.org/10.3168/jds.2015-10012>.
- Li, B., Fikse, W.F., Lassen, J., Lidauer, M.H., Løvendahl, P., Mäntysaari, P. and Berglund, B. 2016. Genetic parameters for dry matter intake in primiparous Holstein, Nordic Red, and Jersey cows in the first half of lactation. *J. Dairy Sci.* 99 (9), 7232-7239. Doi: <https://doi.org/10.3168/jds.2015-10669>.
- López-Paredes, J., Saborío-Montero, A., Charfeddine, N., Jiménez-Montero, J.A., and González-Recio, O. 2021. Dry matter intake, methane emissions and microbiome profiles as new traits for feed efficiency. *Interbull Bulletin* 56, 111-120.
<https://journal.interbull.org/index.php/ib/article/view/77/77>.
- Madsen, P. and Jensen, J. 2013. DMU Ver. 6, rel 5.2.
https://dmu.agrsci.dk/DMU/Doc/Current/dmuv6_guide.5.2.pdf.
- Manzanilla-Pech, C.I.V., Løvendahl, P., Mansan Gordo, D., Difford, G.F., Pryce, J.E., Schenkel, F., Wegmann, S., Miglior, F., Chud, T.C., Moate, P.J., Williams, S.R.O., Richardson, C.M., Stothard, P., and Lassen, J. 2021. Breeding for reduced methane emission and feed-efficient Holstein cows: An international response. *J. Dairy Sci.* 104 (8), 8983-9001.
Doi: <https://doi.org/10.3168/jds.2020-19889>
- Manzanilla-Pech, C.I.V., Stephansen R.B., Difford G.F., Løvendahl P., and Lassen J. 2022. Selecting for feed efficient cows will help to reduce methane gas emissions. *Front. Genet.* (13). Doi: <https://doi.org/10.3389/fgene.2022.885932>
- Manzanilla-Pech, C.I.V., Stephansen, R.B., and Lassen, J. 2023. Genetic parameters for feed intake and body weight in dairy cattle using high-throughput 3-dimensional cameras in Danish commercial farms. *J. Dairy Sci.* 106 (12), 9006-9015.
Doi: <https://doi.org/10.3168/jds.2023-23405>
- Stephansen, R.B., Lidauer, M.H., Nielsen, S., Pösö, J., Fikse, F., Manzanilla-Pech, C.I.V., and Aamand, G.P. 2021. Genomic prediction of residual feed intake in the Nordic breeds using data from research herds and 3D cameras in commercial herds. *Interbull Bulletin* 56, 166-162.
<https://journal.interbull.org/index.php/ib/article/view/66/66>.
- Toshniwal, J.K., Dechow, C.D., Cassell, B.G., Appuhamy, J.A.D.R.N., and Varga, G.A. 2008. Heritability of electronically recorded daily body weight and correlations with yield, dry matter intake, and body condition score. *J. Dairy Sci* 91 (8), 3201–3210. Doi: <https://doi.org/10.3168/jds.2007-0627>.
- Viking Genetics. 2021. Feed Efficiency takes a major step forward. Assessed 05.06.2024.
<https://www.vikinggenetics.com/news/feed-efficiency-takes-a-major-step-forward>.
- Wallén, S.E., Prestløkken, E., Meuwissen, T.H.E., McParland, S., and Berry, D.P. 2018. Milk mid-infrared spectral data as a tool to predict feed intake in lactating Norwegian Red dairy cows. *J. Dairy Sci* 101 (7), 6232-6243. Doi: <https://doi.org/10.3168/jds.2017-13874>.
- Wethal, K.B., Difford, G.F., Winnberg, K., Norberg, E., and Heringstad, B. 2022. Heritability of methane emissions in Norwegian red cows based on measures from GreenFeed in commercial herds. Proc. 12th World Congress on Genetics Applied to Livestock Production, Rotterdam, the Netherlands, Wageningen Academic Publishers, pp. 176-178.